

# Metadata

## Dataset Title:

Cary\_Met\_Data\_3min.csv

Cary Environmental Monitoring Program 3-Minute Meteorological and Solar Radiation Data: 2011-2018

## Abstract

The Cary Institute of Ecosystem Studies Environmental Monitoring Program is a long-term data collection program designed to understand how the environment changes over time. The program includes monitoring of climate including temperature and precipitation, as well as variables related to air pollution, such as acid deposition and ozone, and water pollution and other streamwater chemistry. Our solar radiation monitoring includes diffuse and global photosynthetically active radiation (PAR), diffuse and global shortwave radiation, net radiation and UV. Long-term monitoring of solar radiation provides us with an understanding of atmospheric energy dynamics, which can affect natural and human systems. The Cary Institute of Ecosystem Studies, Environmental Monitoring Program furnishes data under the following conditions: The data have received quality assurance scrutiny by our program, and, although we are confident of the accuracy of these data, the Cary Institute will not be held liable for errors in these data. Data are subject to change resulting from updates in data screening or models used. Data citation: The following is a standard citation for referencing data from the Cary Institute of Ecosystem Studies, Environmental Monitoring Program:

Cary Institute of Ecosystem Studies, Environmental Monitoring Program. 2008 (or current year). Cary Institute of Ecosystem Studies, Box AB, Millbrook, NY 12545, [www.caryinstitute.org](http://www.caryinstitute.org).

Those wishing to publish data from the Cary Institute of Ecosystem Studies, Environmental Monitoring Program are encouraged to contact Data Manager Vicky Kelly, [kellyv@caryinstitute.org](mailto:kellyv@caryinstitute.org).

## Investigators

| First Name | Middle Initial | Last Name | Organization                        | e-mail address   | ORCID ID            |
|------------|----------------|-----------|-------------------------------------|--|---------------------|
| Victoria   | R              | Kelly     | Cary Institute of Ecosystem Studies | <a href="mailto:kellyv@caryinstitute.org">kellyv@caryinstitute.org</a> | 0000-0003-3418-9211 |

## Keywords

Cary Institute, meteorology, climate, solar radiation, PAR, net radiation, shortwave radiation, air temperature, precipitation, barometric pressure, wind speed, wind direction, relative humidity

## Timeframe

The data in this file start 14 July 2011 and end 31 December 2018. Data collection is ongoing.

## Geographic location

The meteorological and solar radiation instruments are located in a flat, open field in Millbrook, NY at an elevation of 128 m. GPS coordinates are N41.785823 W073.741447.

## Methods

Meteorological and solar radiation data are collected using a Campbell Scientific, Inc. datalogger (21X 1M 1987-14 September 1993, 21XL 14 September 1993- 12 October 1999, 23X 12 Oct. 1999-December 2011, CR3000 Dec. 2011-present). All sensors are sampled every 2 seconds and all data are summarized every 3,

minutes, 15 minutes, hourly and at the end of every 24-hour period. Hour-minute data units are Eastern Standard Time (EST), where 1115 is 11:12 to 11:15 EST. Three minute summary meteorological and solar radiation data have been collected since 2011 and 15-minute data since 2015. The meteorological and solar radiation instruments are located in a flat, open field at an elevation of 128 m. GPS coordinates for the site are: N41.785823 W073.741447.

The instruments, their accuracy and our recommendation for the number of significant figures to use:

Please note that significant figures may be reported beyond the accuracy of the instruments. Data are reported in this manner to maintain the capability of backing out calculations from the raw measurements. Instrument notes including calibration, malfunctions, repairs and anecdotal information is available upon request. The instruments, their accuracy and our recommendation for the number of significant figures to use are:

Barometric Pressure, October 1999 - present Campbell Scientific, Inc. Model CS105, sensitivity + 4 mb, recommended significant figs. 4. The barometric pressure sensor is located adjacent to the datalogger in a temperature-controlled room.

Net Radiation, 1988 - April 1992, Qualimetrics Fritschen Type Model 3030 3032, sensitivity + 4 mV, recommended significant figs. 2. Net Radiation, April 1992 - March 1994, Radiation and Energy Balance Systems (REBS) Model Q6, March 1994-Sept. 1995, REBS Model Q\*6.7, Sept. 1995-present REBS Model Q\*7.1, ventilator used Sept 1995-July 2000, sensitivity 0.01 Wm<sup>-2</sup>-mV, recommended significant figs. 2. Location of sensors: 2 m above a mowed grass surface (1988-July 2002), 2.5 m above mowed grass surface (July 2002-present).

PAR, 1988 - present, Li-Cor, Inc. Model LI-190SB, sensitivity 8 mA/1000 mmols-1-m<sup>2</sup>, recommended significant figs. 3. Location of sensors: 2 m above a mowed grass surface (1988-July 2002), 2.5 m above mowed grass surface (July 2002-present).

Relative Humidity, 1988 - April 1997, Phys Chem Corp. PCRC-11 or PCRC-55, sensitivity + 3%, recommended significant figs. 2. Relative Humidity, April 1997-present, Campbell Scientific, Inc. HMP45C, which uses a Vaisala, Inc., capacitive polymer H chip, sensitivity + 3%, recommended significant figs. 2. Temperature and relative humidity sensors are housed in a motor-aspirated shield (Climatronics Corp. series TS-10) 1.6 m above a mowed grass surface.

Shortwave Radiation, 1988 - present, Eppley Laboratory, Inc. Model 8-48, sensitivity 11 mV/Wm<sup>-2</sup>, recommended significant figs. 3. Location of sensors: 2 m above a mowed grass surface (1988-July 2002), 2.5 m above mowed grass surface (July 2002-present).

Temperature, 1988 - Nov. 1998, Campbell Scientific Model 107 or 207, sensitivity + 0.4 deg C, recommended significant figs. 3. Temperature, Nov. 1998 - present, Campbell Scientific, Inc. HMP45C temperature probe, sensitivity + 0.4 deg C, recommended significant figs. 3. Temperature and relative humidity sensors are housed in a motor-aspirated shield (Climatronics Corp. series TS-10) 1.6 m above a mowed grass surface.

Wind Speed, 1988 - July 2002, Campbell Scientific, Inc. Model 014A, sensitivity +1.5%, recommended significant figs. 3. Wind Speed, July 2002 - June 2003, Met One Instruments, Inc. Model 50.5 Solid State Wind Sensor, sensitivity +2%, recommended significant figs. 3. Wind Speed, October 2003 - present, Climatronics Corp. sonimometer. Location of sensors: approximately 10 m above a mowed grass surface.

Wind Direction, 1988 - July 2002, Campbell Scientific, Inc. Model 024A, sensitivity +5 degrees, recommended significant figs. 2. Wind Direction, July 2002 - June 2003, Met One Instruments, Inc. Model 50.5 Solid State Wind Sensor, sensitivity +3 degrees, recommended significant figs. 3. Wind Speed, October 2003 - present, Climatronics Corp. sonimometer. Location of sensors: approximately 10 m above a mowed grass surface.

## METEOROLOGICAL DATA QUALITY ASSURANCE AND QUALITY CONTROL PARAMETERS AND METHODS

The quality of meteorological data collected as part of the Cary Institute Environmental Monitoring

Program is assured and controlled at several levels. Instruments are returned to manufacturers for audits and calibration as recommended by the manufacturer or as needed. If an instrument is known to have malfunctioned or if it is not in place for any reason, data are either removed from the database and replaced with missing values, or are replaced with data from a known working instrument.

Data are checked for accuracy at several levels. Meteorological data are collected using a Campbell Scientific datalogger. The datalogger assigns out-of-range data a value of either NAN, 6999, 9999 or -9999. The program used to import data contains range-checking steps. These steps produce a temporary dataset that put out-of-range data and data that are beyond a reasonable range equal to missing values. These data are then checked and missing values are reviewed. Line graphs of raw data are compared with data from previous years to check for data drift and outliers. All outliers are checked and suspect values are replaced with a missing value code.

Instrument notes including calibration schedule, malfunctions and repairs, new instrumentation, anecdotal information etc. can be made available on request.

The quality of meteorological data collected as part of the Cary Institute Environmental Monitoring Program is assured and controlled at several levels. Instruments are returned to manufacturers for audits and calibration as recommended by the manufacturer or as needed. If an instrument is known to have malfunctioned or if it is not in place for any reason, data are either removed from the database and replaced with missing values, or are replaced with data from a known working instrument.

## Data Table

| Column name | Description   | Unit or code explanation or date format | Empty value code               |
|-------------|---|---|--------------------------------|
| DATE        | Date in MM/DD/YYYY format where MM is month number, DD is date number and YYYY is year  | Date                                    |                                |
| HOUR_MIN    | Hour in hour_minute format in Eastern Standard Time. Note that data are from the preceding hour. For example, data for hour 0100 are from midnight to 1:00 a.m EST. | numeric                                 | Eastern Standard Time          |
| MAX_WS      | Maximum relative humidity (%)   | numeric                                 | percent                        |
| AVE_SWS     | Scalar mean wind speed (m/sec)  | numeric                                 | meters per sec                 |
| AVE_RWS     | Resultant mean wind speed (m/sec)   | numeric                                 | meters per sec                 |
| AVE_RWD     | Resultant mean wind direction (degrees)   | numeric                                 | degrees                        |
| WD_SD       | Standard deviation of wind direction (degrees)  | numeric                                 | degrees                        |
| AVE_TEMP    | Average air temperature (degrees C)   | numeric                                 | degrees C                      |
| MAX_RH      | Maximum relative humidity (%)   | numeric                                 | percent                        |
| MIN_RH      | Minimum relative humidity (%)   | numeric                                 | percent                        |
| AVE_GSW     | Average global short-wave radiation (Watts/m <sup>2</sup> )   | numeric                                 | Watts/m <sup>2</sup>           |
| AVE_DSW     | Average diffuse short-wave radiation (Watts/m <sup>2</sup> )  | numeric                                 | Watts/m <sup>2</sup>           |
| AVE_GPAR    | Average global PAR (micromoles/m <sup>2</sup> -sec)   | numeric                                 | micromoles/m <sup>2</sup> -sec |
| AVE_DPAR    | Average diffuse PAR (micromoles/m <sup>2</sup> -sec)  | numeric                                 | micromoles/m <sup>2</sup> -sec |

|          |  |         |                            |
|----------|--|---------|----------------------------|
| AVE_NETR | Average net radiation (Watts/m^2)      | numeric | Watts per<br>meter squared |
| AVE_BP   | Average barometric pressure (mm<br>Hg) | numeric | mm Hg                      |

## Intellectual Rights

This information is released under the Creative Commons license - Attribution-NonCommercial - CC BY-NC (<https://creativecommons.org/licenses/by/4.0/>).